


Single Top Quark at the Tevatron

Cecilia E. Gerber

University of Illinois at Chicago
for the CDF & DØ collaborations




Top 2011 Workshop
Sant Feliu de Guixols
25-30 Sept. 2011




Outline

- Introduction
- Refresher of Observation Analyses
- Details of new 5.4fb^{-1} DØ results
- Understanding the data
 - Event Selection
 - Background Modeling
- Multivariate Analysis Techniques
- Cross Sections
- Direct Measurement of $|V_{tb}|$
- Anomalous Couplings
- Conclusions & Outlook




Why Study Single Top Production?

- Probe of the Wtb interaction with no assumption on the number of quark families or unitarity of the CKM matrix
- Allows for indirect measurement of the top quark width
- Cross sections sensitive to beyond-the-SM processes
 - Resonances
 - heavy W' boson, charged Higgs boson, Kaluza-Klein excited W_{KK} , technipion, etc.
 - flavor-changing neutral currents
 - Fourth generation of quarks



s-channel

$$\sigma_{\text{SM}} = 0.88 \pm 0.11 \text{ pb}$$





t-channel

$$\sigma_{\text{SM}} = 1.98 \pm 0.25 \text{ pb}$$


Single top cross sections from Kidonakis PRD 74, 114012 (2006) for $m_t = 172.5 \text{ GeV}$

Experimentally very challenging





Simple counting experiment cannot extract the signal from the overwhelming background

Experimentally very challenging



Same final state as WH

- Backgrounds are the same
- Test of techniques to extract small signal from a large background




Observation of Single Top Production

March 2009, 14 years after pair production, 50 times more data

Signal Significance	
Expected	Observed
DØ (2.3 fb⁻¹) PRL 103, 092001 (2009)	
4.5 σ	5.0 σ
CDF (3.2 fb⁻¹) PRL 103, 092002 (2009)	
> 5.9 σ	5.0 σ

$$|V_{tb}| = 0.88 \pm 0.07$$

$$|V_{tb}| > 0.77 @ 95\% C.L.$$



Culmination of a Long History

DØ

- Search: PRD 63, 031101 (2000)
- Search: PLB 517, 282 (2001)
- Search: PLB 622, 265 (2005)
- W' : PLB 641, 423 (2006)
- Search: PRD 75, 092007 (2007)
- **Evidence:** PRL 98, 181802 (2007)
- FCNC: PRL 99, 191802 (2007)
- W' : PRL 100, 211802 (2007)
- Evidence: PRD 78, 012005 (2008)
- Wtb: PRL 101, 221801 (2008)
- Wtb: PRL 102, 092002 (2009)
- H^+ : PRL 102, 191802 (2009)
- **Observation:** PRL 103, 092001 (2009)

CDF

- Search: PRD 65, 091102 (2002)
- W' : PRL 90, 081802 (2003)
- Search: PRD 69, 052003 (2004)
- Search: PRD 71, 012005 (2005)
- **Evidence:** PRL 101, 252001 (2008)
- FCNC: PRL 102, 151801 (2009)
- W' : PRL 103 091801 (2009)
- **Observation:** PRL 103, 092002 (2009)

Cecilia E. Gerber (UIC) – TOP2011

PRL 103, 092001 (2009) PHYSICAL REVIEW LETTERS week ending 28 AUGUST 2009

Observation of Single Top-Quark Production

V.M. Abazov,³⁶ B. Abbott,⁷⁴ M. Abolins,⁶⁴ B.S. Acharya,²⁹ M. Adams,⁵⁰ T. Adams,⁴⁸ E. Aguilo,⁶ M. Ahsan,⁵⁸ G.D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,^{64*} G. Alverem,⁶² C. A. Alvarado,² I. S. Anan,³⁵ T. Anderson,⁵² M. S. Anzures,⁵² M. Aoki,⁷⁹ Y. Arnold,¹⁴ M. Arov,⁵⁹ M. Arthaud,¹ J. BackusMayes,⁴¹ F. Badaud,¹³ L. Bagby,⁴⁹ B. Baldin,¹ A.-F. Barfuss,¹⁵ P. Bargassa,⁷⁹ P. Baringer,⁵⁷ J. Barrelet,¹ M. Begalli,³ M. Begel,⁷² C. Belanger-Champagne,⁴¹ G. Bernardi,¹⁷ R. Bernhard,²³ J. Bertram,⁴² M. B. V. Bhatnagar,²⁷ G. Blazey,⁵¹ S. Blessing,⁴⁸ K. Bloom,¹ G. Borisov,⁴² T. Bose,⁷⁶ A. Brandt,⁷⁷ R. Brock,⁶⁴ G. B. D. Buchholz,⁵² M. Buehler,⁸⁰ V. Buescher,²² V. Bunich,¹ B. Calpas,¹⁵ S. Calvet,¹⁶ J. Cammin,⁷⁰ M. A. Carrá,¹ H. Castilla-Valdez,³³ S. Chakrabarti,⁷¹ D. Chakraborty,¹ B. Choudhary,²⁸ L. Christofek,⁷⁶ T. Christoudias,⁴³ S. W. E. Cooper,⁴⁹ M. Corcoran,⁷⁹ F. Coudere,¹⁸ M. C. Cwiok,⁴⁰ A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁷ S. J. de M. Demarteau,⁴⁹ R. Demina,⁷⁰ D. Denisov,⁴⁹ S. P. Den T. Dorland,⁸¹ A. Dubev,²⁸ L. V. Dudko,³⁸ L. Duflot,¹⁶ D. Duggan,¹ A. Duperrin,¹ S. Dutt,¹ A. Dvshkant,¹ M. Eads,¹

Observation papers
PRL “Editor’s Suggestion”
Physics Synopsis Article
TOPCITE 100+

PRL 103, 092002 (2009) PHYSICAL REVIEW LETTERS week ending 28 AUGUST 2009

Observation of Electroweak Single Top-Quark Production

T. Aaltonen,²⁴ J. Adelman,¹⁴ T. Akimoto,⁵⁶ B. Álvarez González,^{12*} S. Amerio,^{44b,44a} D. Amidei,³⁵ A. Anastassov,³⁰ A. Annovi,²⁰ J. Antos,¹⁵ G. Apollinari,¹⁸ A. Apresyan,⁴⁹ T. Arisawa,⁵⁸ A. Artikov,¹⁶ W. Ashmanskas,¹⁸ A. Attal,⁴

Papers Since Observation

DØ

- t-channel: PLB 682, 81 (2010)
- Tau+jets: PLB 690, 05 (2010)
- FCNC: PLB 693, 81 (2010)
- W' : PLB 699, 145 (2011)
- Top Width: PRL 106, 022001 (2011)

CDF






- MET + Jets: PRD81, 072003 (2010)
- Observation: PRD 82, 112005 (2010)

A. Auris,¹ V. Bartsch,³¹ D. Benjamin,¹ C. Blocke,¹ A. Boveia,¹¹ S. Budd,¹ C. Calancha,¹ S. Carrillo,¹⁹ M. Cavalli-Sforza,¹ F. Chlebana,¹ Y. S. Ci,¹ M. E. C,¹ C. Cuen,¹ P. de Barbaro,¹ M. Deninno,⁶ M. D’Onofri,¹ D. Errede,¹ J. P. Fernandez,¹ I. Furic,¹⁹ M. H. Gerb,¹ J. L. Gim,¹ V. Glagolev,¹ M. Goncl,¹ C. Grosso-Pil,¹ S. R. Hahn,¹⁴ R. F. Har,¹ C. Henderson,¹ S. Hou,¹ J. Incandella,¹ S. Jindaria,¹ P. E. Karch,¹ H. W. Kim,²⁸ B. Knuteson,¹ D. Krop,¹⁴ N. Krumnack,² M. Kruse,¹ V. Krutelyov,¹ T. Kubo,²⁰ T. Kuhr,⁴¹ N. P. Kulkarni,¹ M. Kurata,²⁰ S. Kwang,¹ A. T. Laasanen,⁴⁹ S. Lami,^{47a} S. Lamm,¹⁸ M. Lancaster,³¹ R. L. Lander,¹ K. Lannon,^{40a} A. Lath,⁵³ G. Latino,^{47c,47a}

Variety of Published Results

- Different Final States

- Electron/Muon + Jets
- Tau + jets (DØ)
- MET + jets (CDF)
- Crucial for all the channels
 - b-jets identification




- S:B ~ 1:20 at best

- Uncertainties in background larger than expected signal



- Use MVAs

- BNN, BDT, ME, NEAT
- Both CDF and DØ used a second MVA to increase sensitivity



- Bayesian posterior probability density to extract cross section



MVA Outputs





arXiv:1101.1275



2D measurements (2010)

Phys. Rev. D82,112005 (2010)


PLB 682, 363 (2010)



- Simultaneous measurements of s- and t-channel cross sections.
- No standard model cross section is assumed for either of the signals.

New 5.4fb^{-1} DØ analyses as a detailed example


The Fermilab Tevatron



- proton anti-proton collider
 - Until recently, only place where top quarks had been studied
- Data delivered $\sim 12 \text{ fb}^{-1}$
 - Recorded $\sim 10.5 \text{ fb}^{-1}$

Run II Integrated Luminosity

19 April 2002 - 28 August 2011





Results based on $\sim 5.5 \text{ fb}^{-1}$

	Run I	Run IIa	Run IIb
Bunches in Turn	6 × 6	36 × 36	36 × 36
$\sqrt{s}$ (TeV)	1.8	1.96	1.96
Peak L ($\text{cm}^{-2}\text{s}^{-1}$)	1.6×10^{30}	9×10^{31}	3×10^{32}
$\int \text{Ldt}$ ($\text{pb}^{-1}/\text{week}$)	3	17	50
Bunch crossing (ns)	3500	396	396
Interactions/ crossing	2.5	2.3	8

Event Selection

FINAL STATE CONTAINS:

- One high- p_T isolated electron or muon
 - $p_T > 15 \text{ GeV}$, $|\eta| < 1.1(\text{e}) < 2.0(\mu)$
- Large missing transverse energy
 - $E_T^{\text{miss}} > 20 \text{ GeV}$
- 2, 3 or 4 jets
 - $p_T > 25 \text{ GeV}$ (jet1), $> 15 \text{ GeV}$ (other jets)
 - $|\eta| < 3.4$
- Total Transverse Energy and Triangular cuts
 - $H_T > 110 - 160 \text{ GeV}$
 - $|\Delta\phi(\text{jet1}, \text{ME}_T)| \text{ vs } \text{ME}_T$
 - $|\Delta\phi(\text{lepton}, \text{ME}_T)| \text{ vs } \text{ME}_T$
- At least one b-tagged jet



Signal acceptance



$$tb = (2.9 \pm 0.4) \%$$

$$tqb = (2.0 \pm 0.3) \%$$

Signal and Background Models

- Single top quark signals modeled using SINGLETOP

- Based on COMPHEP
- Reproduces NLO kinematic distributions
- PYTHIA for parton hadronization




- Top pair backgrounds modeled using ALPGEN

- PYTHIA for parton hadronization
- Parton-jet matching algorithm used to avoid double-counting final states
- Normalized $\sigma = 7.46 + 0.48 - 0.67$ pb from Moch and Uwer, PRD 78, 034003 (2008) for $m_t = 172.5$ GeV
- Additional uncertainties include color reconnection (1%), jet fragmentation and higher order effects (1.6%-7%), ISR/FSR (0.8%-11%), and b-jet fragmentation (2%).

Signal and Background Models (Cont.)


- W+jets modeled using ALPGEN

- PYTHIA for parton hadronization
- MLM parton-jet matching avoids double-counting final states
- $\eta(\text{jets})$, $\Delta\phi(\text{jet1}, \text{jet2})$, $\Delta\eta(\text{jet1}, \text{jet2})$ corrected to match data



- QCD Multijet

- Misidentified lepton, directly from data
- Kept small ($\sim 5\%$) with topological selection cuts



- Z+jets modeled using ALPGEN + PYTHIA

- Normalized to NNLO given by FEWZ, with 3.3% uncertainty

- Dibosons modeled using PYTHIA



- Normalized to NLO given by MCFM, with 7% uncertainty

Background Normalization

- W+jets and multijet normalized using iterative template fits to data **BEFORE TAGGING** on three sensitive variables: $p_T(\ell)$, E_T , $M_T(W)$

$$N_{\text{pretag}}^{\text{data}} - N_{\text{non W+jets}}^{\text{MC}} = \lambda_{\text{W+jets}} N_{\text{W+jets}}^{\text{MC}} + \lambda_{\text{multijet}} N_{\text{multijet}}^{\text{data}}$$


- $\lambda_{\text{W+jets}}$ & $\lambda_{\text{multijet}}$ are varied to maximize the product of the KS values for the three sensitive variables.
- Uncertainties are 30%-40% for multijet, 1.8% for W 's



S:B = 1:224

b-Jet Identification

- Separate b-jets from light-quark and gluon jets to reject most W +jets background
- DØ uses a neural network algorithm with seven input variables based on impact parameter and reconstructed vertex
- Two operating points used in analysis:
 - TIGHT ($\epsilon_b = 40\%$, $\epsilon_c = 9\%$, $\epsilon_l = 0.4\%$)
 - LOOSE ($\epsilon_b = 50\%$, $\epsilon_c = 14\%$, $\epsilon_l = 1.5\%$)
- Leading b-jet $p_T > 20$ GeV
- Define two exclusive samples
 - EqOneTag: 1T, no L
 - EqTwoTag: 2L
- b-tagging uncertainties dominated by variation in data samples used to measure the efficiencies.
- Smaller contribution from MC sample dependence





W + Heavy Flavor Normalization

- W/Z + heavy flavor normalized to theory (MCFM-NLO)
 - 1.47 (Wbb,Wcc), 1.32 (Wcj), 1.52 (Zbb), 1.67 (Zcc)
- Normalization checked in 2 jets/0 tag sample

$$N^{(0)} = N_{Wlp}^{(0)} + \lambda_{HF} N_{Whp}^{(0)}$$

- λ_{HF} found to be consistent with 1
- Uncertainties considered
 - $\pm 40\%$ single top cross section
 - $\pm 1\%$ in λ_{HF}
 - $\pm 10\%$ on the Wcj theory SF
 - $\pm 7\%$ in λ_{HF}
 - Additional $\pm 10\%$ Wbb/Wcc
 - $\pm 8\%$ in λ_{HF}
 - For a total uncertainty of 12%



Systematic Uncertainties

Relative Systematic Uncertainties

Components for Normalization

Integrated luminosity	6.1%
$t\bar{t}$ cross section	9.0%
Z+jets cross section	3.3%
Diboson cross sections	7.0%
Branching fractions	1.5%
Parton distribution functions (signal acceptances only)	2.0%
Triggers	5.0%
Instantaneous luminosity reweighting	1.0%
Primary vertex selection	1.4%
Color reconnection	1.0%
b/light jet response	(0.3-1.0)%
Electron identification	(2.8-3.8)%
Muon identification	2.1%
Jet fragmentation and higher order effects	(0.7-7.0)%
Initial-and final-state radiation	(0.8-10.9)%
b-jet fragmentation	2.0%
Taggability	(3.1-21.5)%
W+jets heavy-flavor correction	12.0%
Z+jets heavy-flavor correction	12.0%
W+jets normalization to data	1.8%
Multijets normalization to data	(30-40)%
MC and multijets statistics	(0.2-16)%

Components for Normalization and Shape





Jet reconstruction and identification	(0.04-3.7)%
Jet energy resolution	(0.2-11.6)%
Jet energy scale	(0.3-14.6)%
Vertex confirmation	(0.1-9.6)%
b tagging, single-tagged	(4.3-14.0)%
b tagging, double-tagged	(5.8-11.2)%
Angular correction	0.3%

Components that most affect the cross section measurement are shown in *red*

Other important contributions are shown in *pink*

Event yields & Background Modeling

Event yields in 5.4/fb $D\phi$ data	
e, μ , 2,3,4-jets 1,2-tags combined	
<i>t</i> -channel	239 ± 28
s-channel	160 ± 27
W+jets	4943 ± 598
Z+jet, dibosons	576 ± 113
<i>tt</i>	2124 ± 383
Multijets	451 ± 56
Total prediction	8492 ± 987
Data	8471 ± 92




S:B = 1:22 in 1Tag
S:B = 1:16 in 2Tag

Cross Check Samples

- Selected to test background model in regions dominated by one type of background: W+jets or Top Pairs


W + JETS

- 2 jets, 1 *b*-tagged jet
- $H_T(\ell, E_T, \text{allJets}) < 175 \text{ GeV}$



Top Pairs

- 4 jets, 1 or 2 *b*-tagged jets
- $H_T(\ell, E_T, \text{allJets}) > 300 \text{ GeV}$







Analysis Strategy


- Combine variables with large discriminating power into discriminant functions that separate signal from background
 - Boosted Decision Tree (BDT)
 - Bayesian Neural Network (BNN)
 - Neuroevolution of Augmented Topologies (NEAT)
- Correlation between methods $\sim 70\%$, gain by combining them

Individual variables

Preliminary



Yield [Events/0.04]









Multivariate Discriminant Outputs

tb discriminant


tqb discriminant

tb+tqb discriminant



Cross Section Measurement







- Cross Sections are measured by building a Bayesian posterior probability density
- For each analysis, the single top cross section is given by the position of the posterior density peak, with 68% asymmetric interval as uncertainty






- Gaussian prior for systematic uncertainties
 - Correlations of uncertainties properly taken into account
- Flat prior in signal cross sections

Linearity Tests

Use ensembles of pseudo-data to test validity of x-sec extraction method




Measured Cross Sections



m_t	170 GeV	172.5 GeV	175 GeV
tb	$1.20^{+0.62}_{-0.56}$	$0.68^{+0.38}_{-0.35}$	$0.53^{+0.36}_{-0.34}$
SM	$1.12^{+0.04}_{-0.04}$	$1.04^{+0.04}_{-0.04}$	$0.98^{+0.04}_{-0.04}$
tqb	$2.65^{+0.65}_{-0.59}$	$2.86^{+0.69}_{-0.63}$	$2.45^{+0.60}_{-0.57}$
SM	$2.34^{+0.12}_{-0.12}$	$2.26^{+0.12}_{-0.12}$	$2.16^{+0.12}_{-0.12}$
$tb+tqb$	$3.70^{+0.78}_{-0.80}$	$3.43^{+0.73}_{-0.74}$	$2.56^{+0.69}_{-0.61}$
SM	$3.46^{+0.16}_{-0.16}$	$3.30^{+0.16}_{-0.16}$	$3.14^{+0.16}_{-0.16}$


[arXiv:1108.3091](#) submitted to Phys. Rev. D

Distributions for $S:B > 0.24$



CKM Matrix Element V_{tb}

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = V_{CKM} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \mathbf{V_{tb}} \end{pmatrix}$$



- Weak interaction eigenstates and mass eigenstates are not the same: there is mixing between quarks, described by CKM matrix
- General form of the Wtb vertex

$$\Gamma_{Wtb}^{\mu} = -\frac{g}{\sqrt{2}} \mathbf{V_{tb}} \left\{ \gamma^{\mu} [f_1^L P_L + f_1^R P_R] - \frac{i\sigma^{\mu\nu}}{M_W} (p_t - p_b)_{\nu} [f_2^L P_L + f_2^R P_R] \right\}$$


- Measurement assumes SM production mechanisms
 - Pure V–A and CP-conserving interaction ($f_1^R = f_2^L = f_2^R = 0$)
 - f_1^L : strength of the left-handed Wtb coupling, is allowed to be anomalous
 - $|V_{td}|^2 + |V_{ts}|^2 \ll |V_{tb}|^2$
- Does not assume 3 generations or unitarity of the CKM matrix

Measurement of $|V_{tb}|$

- Use the measurement of the $tb+tbq$ cross section to make a direct measurement of $|V_{tb}|$:
 - Calculate a posterior in $|V_{tb}|^2$ $\sigma(tb, tqb) \propto |V_{tb}|^2$
 - Measure the strength of the V–A






$$|V_{tb} f_1^L| = 1.02^{+0.10}_{-0.11}$$



[arXiv:1108.3091](#) submitted to Phys. Rev. D

Model-independent t-channel Cross Section

- Construct a 2D posterior probability density for $t\bar{q}b$ vs $t\bar{b}$ cross sections
 - No constraint is imposed on the relative rates of $t\bar{b}$ and $t\bar{q}b$ production
- Extract the $t\bar{q}b$ cross section from a 1D posterior obtained by integrating over the $t\bar{b}$ axis, with no assumption on the $t\bar{b}$ rate



Observed
significance
> 5SD

[arXiv:1105.2788](https://arxiv.org/abs/1105.2788) submitted to Phys Lett B

Anomalous Wtb Couplings

- Most general, lowest dimension Wtb vertex:

$$\mathcal{L} = -\frac{g}{\sqrt{2}}\bar{b}\gamma^\mu(L_V P_L + R_V P_R)tW_\mu^- - \frac{g}{\sqrt{2}}\bar{b}\frac{i\sigma^{\mu\nu}q_\nu}{M_W}(L_T P_L + R_T P_R)tW_\mu^- + h.c.$$



Left-handed vector Right-handed vector Left-handed tensor Right-handed tensor
 $L_V = V_{tb} \cdot f_{L_V}$ $R_V = V_{tb} \cdot f_{R_V}$ $L_T = V_{tb} \cdot f_{L_T}$ $R_T = V_{tb} \cdot f_{R_T}$

SM : $L_V \equiv |V_{tb}| \simeq 1$ and $R_V = L_T = R_T = 0$

- Assumptions:


- Production (SM & Anom.) only via W boson exchange
- $|V_{td}|^2 + |V_{ts}|^2 \ll |V_{tb}|^2$
- CP-conserving Wtb vertex
- Anomalous couplings in both production and decay

- Event kinematics and angular distributions sensitive to anomalous couplings






Analysis Strategy: Three Scenarios

- Pair the L_V coupling with one non-SM coupling and assume the other two are negligible
 - (L_V, L_T) ; $L_V=L_T=1$; $R_V=R_T=0$
 - includes the interference L_V+L_T
 - (L_V, R_V) ; $L_V=R_V=1$; $L_T=R_T=0$
 - (L_V, R_T) ; $L_V=R_T=1$; $R_V=L_T=0$
- Use BNN. For each Scenario:
 - Signal: one of $L_T/R_V/R_T$
 - Background includes SM L_V
- Additional systematics:
 - tb (3.8%), tqb (5.3%), mixed couplings in prod and decay (15%)



- Build 2D posterior of $|V_{tb} f_{LV}|^2$ vs $|V_{tb} f_X|^2$
 - Obtain 1D posterior for $|V_{tb} f_X|^2$ by integrating over $|V_{tb} f_{LV}|^2$

Results





No evidence for anomalous couplings, set 95% C.L upper limits


Anomalous couplings	(L_V, L_T)	(L_V, R_V)	(L_V, R_T)
Cross Section	< 1.21 pb	< 2.81 pb	< 0.60 pb
Coupling $ V_{tb} \cdot f_X ^2$	< 0.13	< 0.93	< 0.06

Conclusions and Plans

- Vibrant Single Top Program after 2009 observation
- With half the available luminosity analyzed:
 - New cross section measurements with precision of $\sim 20\%$
 - Observation of t-channel production
 - Measurements and limits on V_{tb}
 - No departures from the SM observed
- Going after s-channel with full dataset



Latest measurements and discoveries



http://www-d0.fnal.gov/Run2Physics/top/top_public_web_pages/top_public.html#singletop

http://www-cdf.fnal.gov/physics/new/top/public_singletop.html

